



Outdoor air dominates burden of disease from indoor exposures

Hänninen, O.; Asikainen, A.; Carrer, P.; Kephelopoulos, S.; de Oliveira Fernandes, E.; Wargocki, Pawel

Published in:

Proceedings of Abstracts 9th International Conference on Air Quality

Publication date:

2014

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Hänninen, O., Asikainen, A., Carrer, P., Kephelopoulos, S., de Oliveira Fernandes, E., & Wargocki, P. (2014). Outdoor air dominates burden of disease from indoor exposures. In T. Mitto, J. Fallmann, U. Mikolajczyk, P. Suppan, V. Singh, & R. S. Sokhi (Eds.), *Proceedings of Abstracts 9th International Conference on Air Quality : Science and Application* (pp. 63-64). University of Hertfordshire.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

OUTDOOR AIR DOMINATES BURDEN OF DISEASE FROM INDOOR EXPOSURES

O. Hänninen (1), A. Asikainen (1), P. Carrer (2), S. Kephapoulos (3), E. de Oliveira Fernandes (4), P. Wargocki (5)

(1) National Institute for Health and Welfare, Finland, POB 95, 70701 Kuopio, Finland; (2) University of Milan, Italy, (3) EC Joint Research Centre, Ispra, Italy; (4) University of Porto, Portugal; (5) Technical University of Denmark

Presenting author email: otto.hanninen@thl.fi

Summary

Both indoor and outdoor sources of air pollution have significant public health impacts in Europe. Based on quantitative modelling of the burden of disease the outdoor sources dominate the impacts by a clear margin.

Introduction

Ambient air pollution has been suggested to be the dominating source of environmental burden of diseases in Europe (Hänninen & Knol, 2011, 2013). However, indoor sources can generate extremely high exposure levels in poorly ventilated conditions. Globally it has been estimated that indoor PM exposures, especially due to combustion of solid fuels, dominate the burden of disease caused by air pollution (Lim et al., 2012). The aim of this work is to compare the magnitude of indoor and outdoor air pollution in European countries and further to evaluate their control potentials using ventilation, filtration and source elimination.

Methods

Burden of disease model developed in the IAIAQ-study (Jantunen et al., 2011) was complemented with a mass-balance component (Hänninen et al., 2004). Exposure data for PM_{2.5}, VOCs, radon, CO, second hand smoke and dampness and mould were collected for 26 European countries (EU27 except Malta). Burden of disease for the baseline at 2010 and for alternative control scenarios were evaluated.

Results and discussion

Total burden of disease from indoor exposures is estimated to be 2.1 million DALY in EU26. The burden is dominated by outdoor PM_{2.5} (62%) followed by indoor PM_{2.5}, radon (8%) and dampness and mould (5%). PM_{2.5} burden is further dominated by cardiovascular diseases followed by respiratory diseases.

Outdoor air burden can be controlled by either limiting ventilation or by filtration. However, the former approach increases the burden from indoor sources if they are not specifically controlled first. Most efficient reduction can be achieved by combining these strategies, providing an almost 50% reduction. As the energy efficiency demands are shaping the future building stocks strongly, it is essential to account for also for the indoor source control to protect public health.

Conclusions

Outdoor air remains the dominating source of burden of disease from indoor exposures. Improvement of outdoor air quality has the largest potential for public health improvements.

Acknowledgement

This work was funded by the EU Health Programme projects HEALTHVENT, Grant Nr. 2009 12 08, and IAIAQ, Grant Nr. 2009 62 02, EU FP6 project EnVIE, SSPE-CT-2004-502671, and Ministry of Social Affairs and Health and intramural funding for project EBoDE.

References

Hänninen O, Knol A (ed.), 2011. European perspectives on Environmental Burden of Disease; Estimates for nine stressors in six countries. THL Reports 1/2011, Helsinki, Finland. 86 pp + 2 appendixes. ISBN 978-952-245-413-3. <http://www.thl.fi/thl-client/pdfs/b75f6999-e7c4-4550-a939-3bccb19e41c1> (accessed 2011-03-23).

Hänninen, O, Knol A, Jantunen M, Lim T-A, Conrad A, Rappolder M, Carrer P, Fanetti A-C, Kim R, Buekers J, Torfs R, Iavarone I, Classen T, Hornberg C, Mekel O, and the EBoDE Group, 2013. Environmental Burden of Disease in Europe: estimates for nine stressors in six countries. Environmental Health Perspectives: Provisionally accepted.

Hänninen, O.O., Lebrecht, E., Ilacqua, V., Katsouyanni, K., Künzli, N., Srám, R.J., Jantunen, M.J., 2004. Infiltration of ambient PM_{2.5} and levels of indoor generated non-ETS PM_{2.5} in residences of four European cities. Atmospheric Environment 38 (37): 6411-6423.

Jantunen M, Oliveira Fernandes E, Carrer P, Kephapoulos S, 2011. Promoting actions for healthy indoor air (IAIAQ). European Commission Directorate General for Health and Consumers. Luxembourg. ISBN 978-92-79-20419-7. http://ec.europa.eu/health/healthy_environments/docs/env_iaiaq.pdf (accessed 2011-11-10)